

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

AMENDMENTS TO THE CLAIMS

D²
1. (Currently Amended) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material, comprising the step of adding a structured silicate salt ~~in which the cation is~~ contains a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof to a binder of an electrophotographic toner or developer or of an electret material.

2) (Cancelled)*









3) (Cancelled)

4) (Original) The method as claimed in claim 1, wherein the low molecular weight organic cation is a substituted ammonium, phosphonium, thionium or triphenylcarbonium ion or a cationic metal complex.

D³
5) (Currently Amended) The method as claimed in claim 4, wherein the ammonium ion has one of the formulae (a) - (j)



DESI AVAILABLE COPY

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756



BEST AVAILABLE COPY


Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756



in which

R^1 to R^{18} are identical or different and represent hydrogen, CN, $(CH_2)_{1-18}$ CN, halogen, branched or unbranched C_1 - C_{32} -alkyl, mono- or polyunsaturated C_2 - C_{32} -alkenyl, C_1 - C_{22} -alkoxy, C_1 - C_{22} -hydroxyalkyl, C_1 - C_{22} -halogenoalkyl, C_2 - C_{22} -halogenoalkenyl, C_1 - C_{22} -aminoalkyl, $(C_1$ - C_{12})-trialkyl-ammonium- $(C_1$ - $C_{22})$ -alkyl; $(C_1$ - $C_{22})$ -alkylene-(C=O)O- $(C_1$ - $C_{32})$ -alkyl, $(C_1$ - $C_{22})$ -alkylene-(C=O)O-aryl, $(C_1$ - $C_{22})$ -alkylene-(C=O)NH- $(C_1$ - $C_{32})$ -alkyl, $(C_1$ - $C_{22})$ -alkylene-(C=O)NH-aryl, $(C_1$ - $C_{22})$ -alkylene-O(CO)- $(C_1$ - $C_{32})$ -alkyl, $(C_1$ - $C_{22})$ -alkylene-O(CO)aryl, $(C_1$ - $C_{22})$ -alkylene-NH(C=O)- $(C_1$ - $C_{32})$ -alkyl, $(C_1$ - $C_{22})$ -alkylene-NHCO-aryl,

wherein



are optionally inserted into the acid ester or acid amide bonds;





$[(C_1$ - $C_{12})$ -alkylene-O-] $_{1-100}$ -H; aryl, $(C_1$ - $C_{18})$ -alkylenearyl; $-(O-SiR'_2)_{1-32}-O-SiR'_3$, in which R' has the meaning C_1 - C_{12} -alkyl, phenyl, benzyl or C_1 - C_{12} -alkoxy; heterocyclyl, C_1 - C_{18} -alkylene-heterocyclyl, wherein the aryl and heterocyclyl radicals are optionally mono- or polysubstituted on carbon atoms or heteroatoms by C_1 - C_{12} -alkyl, C_1 - C_4 -alkenyl, C_1 - C_4 -alkoxy, hydroxy- $(C_1$ - $C_4)$ -alkyl, amino- $(C_1$ - $C_4)$ -alkyl, C_1 - C_4 -alkylimino, carboxyl, hydroxyl, amino, nitro, cyano, halogen, C_1 - C_{12} -acyl, C_1 - C_4 -halogenoalkyl, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkylcarbonyloxy, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756

alkylaminocarbonyl, C₁-C₄-alkylcarbonylimino, C₆-C₁₀-arylcarbonyl, aminocarbonyl, aminosulfonyl, C₁-C₄-alkylaminosulfonyl, phenyl, naphthyl, or heteroaryl[2];

R¹⁹ represents C₄-C₁₁-alkylene, -(C₂H₄-O-)₁₋₁₇-(CH₂)₁₋₂-, -(C₂H₄-NR-)₁₋₁₇-(CH₂)₁₋₂-, in which R is hydrogen or C₁-C₁₂-alkyl;

X has the meaning of Y or -CO-CH₂-CO-,



Y has the meaning -C-, -C-, -C-, -(CH₂)₁₋₁₈-,
 $\begin{array}{c} \text{O} \\ \parallel \\ \text{C} \end{array}$ $\begin{array}{c} \text{S} \\ \parallel \\ \text{C} \end{array}$ $\begin{array}{c} \text{NH} \\ \parallel \\ \text{C} \end{array}$

BEST AVAILABLE COPY

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756



or o-, p-, m-(C₆-C₁₄)-arylene or (C₄-C₁₄)-heteroarylene with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S and a combination thereof;

R⁶⁰ represents C₁-C₃₂-acyl, C₁-C₂₂-alkyl, C₂-C₂₂-alkenyl, C₁-C₁₈-alkylene-C₆-C₁₀-aryl, C₁-C₂₂-alkylene-heterocyclyl, C₆-C₁₀-aryl or (C₄-C₁₄)-heteroaryl with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S, and a combination thereof;


R⁶¹ and R⁶⁴ represent -(CH₂)₁₋₁₈-, C₁-C₁₂-alkylene-C₆-C₁₀-arylene, C₆-C₁₀-arylene, C₀-C₁₂-alkylene-heterocyclyl;

Z represents -NH- or -O-;

A₁⁻ and A₃⁻ represent -COO⁻, -SO₃⁻, -OSO₃⁻, -SO₂⁻, -COS⁻ or -CS₂⁻;

A₂ represents -SO₂Na, -SO₃Na, -SO₂H, -SO₃H or hydrogen;

R⁶⁹ and R⁷⁰ independently of one another represent hydrogen, C₁-C₃₂-alkyl, in which the alkyl chain optionally contain one or more of the groups -NH-CO-, -CO-NH-, -CO-O- or -O-CO-; C₁-C₁₈-alkylene-aryl, C₀-C₁₈-alkylene-heterocyclyl, C₁-C₁₈-hydroxyalkyl, C₁-C₁₈-halogenoalkyl, aryl, -(CH₂)₃-SO₃⁻,



Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

13
V
R⁷¹ and R⁷² represent -(CH₂)₁₋₁₂; and
R⁷³ and R⁷⁴ represent hydrogen or C₁-C₂₂-alkyl.

6) (Original) The method as claimed in claim 5, wherein R¹ to R¹⁸ denote hydrogen, CN, CH₂-CN, CF₃, C₁-C₂₂-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₈-alkoxy, C₁-C₁₈-hydroxy-alkyl, C₁-C₁₈-halogenoalkyl, C₂-C₁₈-halogenoalkenyl, C₁-C₁₈-aminoalkyl, (C₁-C₆)-trialkylammonium-(C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkylene-O(C=O)-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-O(C=O)-phenyl, (C₁-C₁₈)-alkylene-NHCO-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-NHCO-phenyl, (C₁-C₁₈)-alkylene-(C=O)O-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-(C=O)O-phenyl, (C₁-C₁₈)-alkylene-(C=O)NH-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-CONH-phenyl, benzyl, phenyl, naphthyl, C₁-C₁₂-alkylene-heterocyclyl;
R¹⁹ denotes C₄-C₅-alkylene, -(C₂H₄-O)₁₋₉-(CH₂)₁₋₂ or -(C₂H₄-NH)₁₋₉-(CH₂)₁₋₂;
R⁶⁰ denotes C₁-C₁₈-acyl, C₁-C₁₈-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₂-alkylene-phenyl, C₁-C₁₈-alkylene-pyridyl, phenyl or pyridyl;
R⁶¹ and R⁶⁴ denote -(CH₂)₁₋₁₂, C₁-C₈-alkylene-phenylene, phenylene or C₁-C₈-alkylenepyridylene or piperidylene;
R⁷¹ and R⁷² denote -(CH₂)₁₋₈ and
R⁷³ and R⁷⁴ denote hydrogen or (C₁-C₁₈)-alkyl.

7) (Previously Presented) The method as claimed in claim 4, wherein the ammonium ion is an aliphatic or aromatic 5- to 12-membered heterocyclic radical with 1 to 4 atoms selected from the group consisting of N, O and S, or a combination thereof, belonging to the rings.


8) (Original) The method as claimed in claim 7, wherein the heterocyclic radical is pyridinium, pyridazinium, pyrimidinium, pyrazinium, purinium, tetraazaporphyrinium, piperidinium, morpholinium, tetrazonium, triaza-cyclononanium or tetraaza-cyclododecanium.

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

9) (Original) The method as claimed in claim 4, wherein the cationic metal complex is a metal carboxylate, metal salicylate, metal sulfonate, 1:1 metal-azo complex or a metal dithiocarbamate.

10) (Previously Presented) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO.

11) (Original) The method as claimed in claim 1, wherein the organic cation is a fluorinated ammonium ion of the formula (x)



in which

R^{28} denotes perfluorinated alkyl having 5 to 11 carbon atoms and

R^{29} , R^{30} and R^{31} are identical or different and denote alkyl having 1 to 5 carbon atoms.

12) (Original) Salt-like structured silicate, in which the silicate is hectorite, beldellite, illite, muscovite, xantophyllite, margarite, sepiolite, saponite, mica, feldspar, nontronite, montmorillonite, smectite, bentonite, faujasite, zeolite A, X or Y, permutite, sasil or a combination thereof; and the cation is an ion of the formula (x) as claimed in claim 9.

Attorney's Docket: 1999DE132
Serial No.: 08/722,760
Group: 1756

13) (Original) A process for the preparation of a salt-like structured silicate as claimed in claim 12, which comprises combining the silicate and a salt of the cation of formula (x) in an aqueous medium.

14) (Currently Amended) An electrophotographic toner comprising 30 to 99.99% by weight of a binder, and 0.01 to 50% by weight, of at least one salt of ionic structured silicates in which the cation is contains a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the electrophotographic toner.

15) (Previously Presented) An electrophotographic toner as claimed in claim 14, comprising 40 to 99.5% by weight of a binder, and 0.05 to 20% by weight of at least one salt of ionic structured silicates in which the cation is a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the electrophotographic toner.

16. (Previously Presented) The method of claim 4, wherein the ammonium ion is an aliphatic or aromatic 5- to 12-membered heterocyclic radical with 1 to 4 atoms selected from the group consisting of N, O and S, or a combination thereof, belonging to the rings, wherein 2 to 8 rings are fused.

17. (Previously Presented) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO, and the metal complex contains one or more further ligands.

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

18. (Currently Amended) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, of a powder coating, or of an electret material, comprising the steps of adding a salt structured silicate ~~in which the cation is or contains~~ a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof to a binder of an electrophotographic toner or developer or of a powder coating, or to an electret material.

19. (Previously Presented) An electrophotographic toner or developer comprising distearyldimethyl ammonium bentonite.

20. (Previously Presented) The electrophotographic toner as claimed in claim 14, further comprising 0.001 to 50% by weight, of a coloring agent, based on the total weight of the electrophotographic toner.

21. (Currently Amended) A composition comprising 30 to 99.99% by weight of a binder, and 0.01 to 50% by weight, of at least one salt of ionic structured silicates in which ~~the cation is~~ contains a low molecular weight organic cation and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof, based on the total weight of the composition, wherein the composition is an electrophotographic toner.

22. (Currently Amended) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material comprising the step of adding a distearyldimethyl ammonium bentonite to a binder of an electrophotographic toner or developer ~~or of a powder coating or of an electret~~ material.